

E_1 -DEGENERATION AND $d'd''$ -LEMMA

TAI-WEI CHEN, CHUNG-I HO, AND JYH-HAUR TEH*

ABSTRACT. For a double complex (A, d', d'') , we show that if it satisfies the $d'd''$ -lemma and the spectral sequence $\{E_r^{p,q}\}$ induced by A does not degenerate at E_0 , then it degenerates at E_1 . We apply this result to prove the degeneration at E_1 of a Hodge-de Rham spectral sequence on compact bi-generalized Hermitian manifolds that satisfy a version of $d'd''$ -lemma.

Keywords: $\partial\bar{\partial}$ -lemma, Hodge-de Rham spectral sequence, E_1 -degeneration, bi-generalized Hermitian manifold.

1. INTRODUCTION

Complex manifolds that satisfy the $\partial\bar{\partial}$ -lemma enjoy some nice properties such as they are formal manifolds ([DGMS]), their Bott-Chern cohomology, Aeppli cohomology and Dolbeault cohomology are all isomorphic. Compact Kähler manifolds are examples of such manifolds. The Hodge-de Rham spectral sequence $E_*^{*,*}$ of a complex manifold M is built from the double complex $(\Omega^{*,*}(M), \partial, \bar{\partial})$ of complex differential forms which relates the Dolbeault cohomology of M to the de Rham cohomology of M . It is well known that $E_1^{p,q}$ is isomorphic to $H^p(M, \Omega^q)$ and the spectral sequence $E_r^{*,*}$ converges to $H^*(M, \mathbb{C})$. The goal of this paper is to prove an algebraic version of the result that the $\partial\bar{\partial}$ -lemma implies the E_1 -degeneration of a Hodge-de Rham spectral sequence. The following is our main result.

Theorem 1.1. *If a double complex (A, d', d'') satisfies the $d'd''$ -lemma and the spectral sequence $\{E_r^{p,q}\}$ induced by A does not degenerate at E_0 , then it degenerates at E_1 .*

We define a spectral sequence that is analogous to the Hodge-de Rham spectral sequence of complex manifolds for bi-generalized Hermitian manifolds. Applying result above, we are able to show that for compact bi-generalized Hermitian manifolds that satisfy a version of $\partial\bar{\partial}$ -lemma, the sequence degenerates at E_1 .

Acknowledgements The authors thank the referee for his/her extremely careful review which largely improves this paper.

2. DEGENERATION OF A HODGE-DE RHAM SPECTRAL SEQUENCE

Definition 2.1. A spectral sequence is a sequence of differential bi-graded modules $\{(E_r^{*,*}, d_r)\}$ such that d_r is of degree $(r, 1-r)$ and $E_{r+1}^{p,q}$ is isomorphic to $H^{p,q}(E_r^{*,*}, d_r)$.

Definition 2.2. A filtered differential graded module is a \mathbb{N} -graded module $A = \bigoplus_{k=0}^{\infty} A^k$, endowed with a filtration F and a linear map $d : A \rightarrow A$ satisfying

- (1) d is of degree 1: $d(A^k) \subset A^{k+1}$;
- (2) $d \circ d = 0$;

*Corresponding author.

2010 *Mathematics Subject Classification.* 55T05, 53C05.

(3) the filtered structure is descending:

$$A = F^0 A \supseteq F^1 A \supseteq \cdots \supseteq F^k A \supseteq F^{k+1} A \supseteq \cdots ;$$

(4) the map d preserves the filtered structure: $d(F^k A) \subset F^k A$ for all k .

For $p, q, r \in \mathbb{Z}$, let

$$\begin{aligned} Z_r^{p,q} &= \left\{ \xi \in F^p A^{p+q} \mid d\xi \in F^{p+r} A^{p+q+1} \right\}, & Z_\infty^{p,q} &= F^p A^{p+q} \cap \ker d \\ B_r^{p,q} &= F^p A^{p+q} \cap dF^{p-r} A^{p+q-1}, & B_\infty^{p,q} &= F^p A^{p+q} \cap \text{Im} d \\ E_r^{p,q} &= \frac{Z_r^{p,q}}{Z_{r-1}^{p+1,q-1} + B_{r-1}^{p,q}}, & E_\infty^{p,q} &= \frac{F^p A^{p+q} \cap \ker d}{F^{p+1} A^{p+q} \cap \ker d + F^p A^{p+q} \cap \text{Im} d} \end{aligned}$$

with the convention $F^{-k} A^{p+q} = A^{p+q}$ and $A^{-k} = \{0\}$ for $k \geq 0$. Let $d_r : E_r^{p,q} \rightarrow E_r^{p+r,q-r+1}$ be the differential induced by $d : Z_r^{p,q} \rightarrow Z_r^{p+r,q-r+1}$.

Throughout this paper, we always assume that $A = \oplus_{p,q \geq 0} A^{p,q}$ is a double complex of vector spaces over some field with two maps $d'_{p,q} : A^{p,q} \rightarrow A^{p+1,q}$ and $d''_{p,q} : A^{p,q} \rightarrow A^{p,q+1}$ satisfying $d'_{p+1,q} d'_{p,q} = 0$, $d''_{p,q+1} d''_{p,q} = 0$ and $d'_{p,q+1} d''_{p,q} + d''_{p+1,q} d'_{p,q} = 0$ for all $p, q \geq 0$. To make notation cleaner, we allow p, q to be any integers by defining $A^{p,q} = 0$ for $p < 0$ or $q < 0$.

Let $A^k = \bigoplus_{p+q=k} A^{p,q}$. Define

$$F^p A^k = \bigoplus_{s=p}^k A^{s,k-s}$$

For $p > k$, define $F^p A^k = \{0\}$. This gives a descending filtration on A^k .

Let $d = d' + d''$. The double complex (A, d', d'') then defines a filtered differential graded module (A, d, F) . Let $\{E_r^{p,q}\}$ be the corresponding spectral sequence. We are interested in the convergence of $E_r^{p,q}$.

Definition 2.3. Let $\{E_r^{p,q}\}$ be the spectral sequence associated to the double complex (A, d', d'') . If $d_s = 0$ for all $s \geq r$, then we say that $\{E_r^{p,q}\}$ or A degenerates at E_r .

The following simple lemmas will be used frequently.

Lemma 2.4. If G' is a vector space and $H < G, H < H'$ are subspaces of G' , the natural map $\varphi : \frac{G}{H} \rightarrow \frac{G'}{H'}$ is injective if and only if $G \cap H' = H$, and is surjective if and only if $G' = G + H'$.

Lemma 2.5. Let $p, q, r \in \mathbb{Z}$. There are inclusions

$$\cdots \subset B_0^{p,q} \subset B_1^{p,q} \subset \cdots \subset B_\infty^{p,q} \subset Z_\infty^{p,q} \subset \cdots \subset Z_1^{p,q} \subset Z_0^{p,q} \subset \cdots ,$$

$$Z_{r-1}^{p+1,q-1} \subset Z_r^{p,q} \quad , \quad B_{r+1}^{p+1,q-1} \subset Z_r^{p,q} \quad , \quad d(Z_r^{p-r,q+r-1}) = B_r^{p,q}$$

Definition 2.6. Let $\alpha_{p,q,r} : E_{r+1}^{p,q} \rightarrow \frac{Z_r^{p,q}}{Z_{r-1}^{p+1,q-1} + B_r^{p,q}}$ be the map induced by the composition of inclusion and projection, and $\beta_{p,q,r} : E_r^{p,q} \rightarrow \frac{Z_r^{p,q}}{Z_{r-1}^{p+1,q-1} + B_r^{p,q}}$ be the map induced by the projection.

Proposition 2.7. Let $r \in \mathbb{Z}$. Then

- (1) $d_r = 0$ if and only if $\beta_{p,q,r}$ is an isomorphism for all $p, q \in \mathbb{Z}$.
- (2) $d_r = 0$ implies that $\alpha_{p,q,r}$ is an isomorphism for all $p, q \in \mathbb{Z}$.

Proof. (1) We first note that the map $\beta_{p,q,r}$ is always surjective. By Lemma 2.4, $\beta_{p,q,r}$ is an isomorphism if and only if $Z_r^{p,q} \cap (Z_{r-1}^{p+1,q-1} + B_r^{p,q}) = Z_{r-1}^{p+1,q-1} + B_r^{p,q}$, or equivalently, $B_r^{p,q} \subseteq Z_{r-1}^{p+1,q-1} + B_{r-1}^{p,q}$. The map $d_r^{p-r,q+r-1} : E_r^{p-r,q+r-1} \rightarrow E_r^{p,q}$ is the zero map if and only if $\text{Im} d_r^{p-r,q+r-1} = \{0\}$. This is equivalent to $d(Z_r^{p-r,q+r-1}) = B_r^{p,q} \subseteq Z_{r-1}^{p+1,q-1} + B_{r-1}^{p,q}$, which is equivalent to $\beta_{p,q,r}$ being an isomorphism.

(2) We recall that the isomorphism $E_{r+1}^{p,q} \xrightarrow{\cong} H^{p,q}(E_r^{*,*}, d_r)$ (see [M, Proof of Theorem 2.6]) is induced from some canonical projections and inclusions. If $d_r = 0$, $H^{p,q}(E_r^{*,*}, d_r) \cong E_r^{p,q}$ and we have a commutative diagram

$$\begin{array}{ccc} E_{r+1}^{p,q} & \xrightarrow{\cong} & E_r^{p,q} \\ \alpha_{p,q,r} \searrow & & \swarrow \beta_{p,q,r} \\ & Z_r^{p,q} & \\ & \underline{Z_{r-1}^{p+1,q-1} + B_r^{p,q}} & \end{array}$$

By (1), $\beta_{p,q,r}$ is an isomorphism and hence $\alpha_{p,q,r}$ is an isomorphism. □

Definition 2.8. Fix a pair of integers (p, q) . For nonzero $\xi = \sum_i \xi_i \in \bigoplus_{i \geq 0} A^{p+i, q-i}$ where $\xi_i \in A^{p+i, q-i}$, let $i_0 = \min_i \{\xi_i \neq 0\}$. We call ξ_{i_0} the leading term of ξ and denote it as $\ell^{p,q}(\xi)$. We define $\ell^{p,q}(0) = 0$. For $r \geq 1, p, q \in \mathbb{Z}$, let

$$\mathcal{E}_r^{p,q} := \left\{ \xi = \xi_0 + \xi_1 + \cdots + \xi_{r-1} \mid \xi_i \in A^{p+i, q-i}, d\xi = d'\xi_{r-1} \notin \text{Im} d'', \ell^{p,q}(\eta) \neq \xi_0 \text{ for all } d\text{-closed } \eta \right\}$$

and

$$\mathcal{E}_0^{p,q-1} := B_0^{p,q} - (Z_{-1}^{p+1,q-1} + B_{-1}^{p,q})$$

Lemma 2.9. Fix $r_0 \geq 1$.

- (1) If the map $\alpha_{p,q,r}$ is an isomorphism for all $p, q \in \mathbb{Z}, r \geq r_0$, then $\mathcal{E}_r^{p,q} = \emptyset$ for all $p, q \in \mathbb{Z}, r \geq r_0$.
- (2) If the map α_{p,q,r_0} is not an isomorphism, then $\mathcal{E}_{r_0}^{p,q} \neq \emptyset$.

Proof. Note that by Lemma 2.4, the surjectivity of $\alpha_{p,q,r}$ is equivalent to the condition

$$Z_r^{p,q} = Z_{r+1}^{p,q} + Z_{r-1}^{p+1,q-1} + B_r^{p,q} = Z_{r+1}^{p,q} + Z_{r-1}^{p+1,q-1}.$$

- (1) Suppose that $\alpha_{p,q,r}$ is an isomorphism for all $r \geq r_0$. Then $Z_i^{p,q} = Z_{i+1}^{p,q} + Z_{i-1}^{p+1,q-1}$ for all $i \geq r_0$. Assume that $\mathcal{E}_r^{p,q} \neq \emptyset$ for some $r \geq r_0, p, q \in \mathbb{Z}$. Let $\xi \in \mathcal{E}_r^{p,q}$. By definition, $Z_{q+2}^{p,q} = Z_{q+3}^{p,q} = \cdots = Z_\infty^{p,q}$. So we may take $j > r$ such that $Z_j^{p,q} = Z_\infty^{p,q}$. Note that $\xi \in Z_r^{p,q}$. Using the relation above, we may write $\xi = \eta_1 + \eta_2$ where $\eta_1 \in Z_j^{p,q}, \eta_2 \in Z_{j-2}^{p+1,q-1} + \cdots + Z_{r-1}^{p+1,q-1}$. Since $\ell^{p,q}(\xi) \neq 0$, by comparing the degrees of both sides of $\xi = \eta_1 + \eta_2$, we have $\ell^{p,q}(\xi) = \ell^{p,q}(\eta_1)$. But $d\eta_1 = 0$ which contradicts to the fact that $\ell^{p,q}(\xi)$ is not the leading term of any d -closed element.

- (2) Fix $r \geq 1$. Suppose that $\alpha_{p,q,r}$ is not an isomorphism, then $Z_{r+1}^{p,q} + Z_{r-1}^{p+1,q-1} \subsetneq Z_r^{p,q}$.

Let

$$\xi = \xi_0 + \xi_1 + \cdots + \xi_k \in Z_r^{p,q} - (Z_{r+1}^{p,q} + Z_{r-1}^{p+1,q-1}) \text{ where } \xi_i \in A^{p+i, q-i}.$$

If $k > r - 1$, let $\xi' = \xi_r + \xi_{r+1} + \cdots + \xi_k \in F^{p+r}A^{p+q} \subset F^{p+1}A^{p+q}$. We have

$$d\xi' = d\xi_r + \cdots + d\xi_k \in F^{p+r}A^{p+q+1} = F^{(p+1)+(r-1)}A^{(p+1)+(q-1)+1}$$

which means that $\xi' \in Z_{r-1}^{p+1,q-1}$. Let $\xi'' = \xi - \xi'$. If $\xi'' \in Z_{r+1}^{p,q} + Z_{r-1}^{p+1,q-1}$, then $\xi = \xi' + \xi'' \in Z_{r+1}^{p,q} + Z_{r-1}^{p+1,q-1}$ which contradicts to our assumption. Therefore $\xi'' = \xi_0 + \cdots + \xi_{r-1} \in Z_r^{p,q} - (Z_{r+1}^{p,q} + Z_{r-1}^{p+1,q-1})$. Hence we may assume $\xi = \xi_0 + \cdots + \xi_{r-1}$.

- (i) Since $\xi \in Z_r^{p,q}$, by definition, $d\xi \in F^{p+r}A^{p+q+1}$. But $d(\xi_0 + \cdots + \xi_{r-2}) + d''\xi_{r-1} \in A^{p,q+1} \oplus A^{p+1,q} \oplus \cdots \oplus A^{p+r-1,q-r+2}$. This forces $d(\xi_0 + \cdots + \xi_{r-2}) + d''\xi_{r-1} = 0$ and hence $d\xi = d'\xi_{r-1}$.
- (ii) If $d'\xi_{r-1} = d''\eta_r$ for some $\eta_r \in A^{p+r,q-r}$, then $d(\xi - \eta_r) = d'\xi_{r-1} - d'\eta_r - d''\eta_r = -d'\eta_r \in A^{p+r+1,q-r} \subset F^{p+(r+1)}A^{p+q+1}$. Hence $\xi - \eta_r \in Z_{r+1}^{p,q}$. Since $\eta_r \in F^pA^{p+q}$ and $d\eta_r \in A^{p+r,q-r+1} \oplus A^{p+r+1,q-r} \subset F^{(p+1)+(r-1)}A^{p+q+1}$, we have $\eta_r \in Z_{r-1}^{p+1,q-1}$. Therefore $\xi = (\xi - \eta_r) + \eta_r \in Z_{r+1}^{p,q} + Z_{r-1}^{p+1,q-1}$ which is a contradiction. Hence $d'\xi_{r-1} \notin \text{Im}d''$.
- (iii) If ξ_0 is the leading term of a d -closed form $\tau \in F^pA^{p+q}$, then $\xi - \tau \in F^{p+1}A^{p+q}$ and $d(\xi - \tau) = d\xi \in F^{p+r}A^{p+q+1} = F^{(p+1)+(r-1)}A^{p+q+1}$. Hence $\xi - \tau \in Z_{r-1}^{p+1,q-1}$. Then $\xi = \tau + (\xi - \tau) \in Z_\infty^{p,q} + Z_{r-1}^{p+1,q-1} \subset Z_{r+1}^{p,q} + Z_{r-1}^{p+1,q-1}$ which is a contradiction. Hence $\xi \in \mathcal{E}_r^{p,q}$.

□

Lemma 2.10. (1) $\mathcal{E}_0^{p,q-1} = \emptyset$ if and only if $\beta_{p,q,0}$ is an isomorphism.

(2) For $r \geq 1$, if $\mathcal{E}_r^{p-r,q+r-1} = \emptyset$, then $\beta_{p,q,r}$ is an isomorphism.

(3) For $r \geq 1$, if $\mathcal{E}_r^{p-r,q+r-1} \neq \emptyset$, then $\beta_{p,q,j}$ is not an isomorphism for $j = 1$ or r .

Proof. We note that $\beta_{p,q,r}$ is an isomorphism if and only if $B_r^{p,q} \subset Z_{r-1}^{p+1,q-1} + B_{r-1}^{p,q}$.

- (1) This follows from the definition.
- (2) Assume that $\beta_{p,q,r}$ is not an isomorphism. Then there exists $\xi \in B_r^{p,q} - (Z_{r-1}^{p+1,q-1} + B_{r-1}^{p,q})$. So $\xi = d\eta$ for some $\eta \in F^{p-r}A^{p+q-1}$. Let

$$\eta = \eta_0 + \eta_1 + \cdots + \eta_k \text{ where } \eta_i \in A^{p-r+i,q+r-i-1}.$$

If $k \geq r$, let $\eta' = \eta_r + \cdots + \eta_k \in F^pA^{p+q-1} \subset F^{p-(r-1)}A^{p+q-1}$. Then $d\eta' \in F^pA^{p+q} \cap d(F^{p-(r-1)}A^{p+q-1}) = B_{r-1}^{p,q}$. If $d(\eta - \eta') \in Z_{r-1}^{p+1,q-1} + B_{r-1}^{p,q}$, then $\xi = d(\eta - \eta') + d\eta' \in Z_{r-1}^{p+1,q-1} + B_{r-1}^{p,q}$ which is a contradiction. So $d(\eta - \eta') \in B_r^{p,q} - (Z_{r-1}^{p+1,q-1} + B_{r-1}^{p,q})$. Hence we may assume $\xi = d\eta$ where $\eta = \eta_0 + \cdots + \eta_{r-1}$.

- (i) Comparing the degrees of ξ and $d\eta$, we see that $d\eta = d'\eta_{r-1}$.
- (ii) If $\eta_0 = 0$, then $\xi = d(\eta_1 + \cdots + \eta_{r-1}) \in F^pA^{p+q} \cap d(F^{p-(r-1)}A^{p+q-1}) = B_{r-1}^{p,q}$ which is a contradiction. So $\eta_0 \neq 0$.
- (iii) If η_0 is the leading term of a d -closed form η'' , $\eta - \eta'' \in F^{p-r+1}A^{p+q-1}$ and $\xi = d\eta = d(\eta - \eta'') \in d(F^{p-(r-1)}A^{p+q-1}) \cap F^pA^{p+q} = B_{r-1}^{p,q}$ which is a contradiction. Hence η_0 is not the leading term of any d -closed form.
- (iv) If $d'\eta_{r-1} \in \text{Im}d''$, $\xi = d\eta = d'\eta_{r-1} = -d''\eta_r$ for some $\eta_r \in A^{p,q-1}$, then $\xi = d'\eta_r - d\eta_r \in Z_\infty^{p+1,q-1} + B_0^{p,q} \subset Z_{r-1}^{p+1,q-1} + B_{r-1}^{p,q}$ which is a contradiction. Hence $d'\eta_{r-1} \notin \text{Im}d''$.

Therefore, $\eta \in \mathcal{E}_r^{p-r,q+r-1}$.

- (3) Assume that $\mathcal{E}_r^{p-r,q+r-1} \neq \emptyset$. Let $\eta = \eta_0 + \cdots + \eta_{r-1} \in \mathcal{E}_r^{p-r,q+r-1}$ where $\eta_i \in A^{p-r+i,q+r-i-1}$. Since $d\eta \in B_r^{p,q}$, if $d\eta \notin Z_{r-1}^{p+1,q-1} + B_{r-1}^{p,q}$, $\beta_{p,q,r}$ is not an isomorphism. So we may assume $d\eta = d'\eta_{r-1} = \xi' + d\eta'$ where $\xi' \in Z_{r-1}^{p+1,q-1}$ and $d\eta' \in B_{r-1}^{p,q}$. Let $\eta' = \eta'_1 + \eta'_2 + \cdots + \eta'_l$, where

$\eta'_i \in A^{p-r+i, q+r-1-i}$. The degree of $d'\eta_{r-1}$ is (p, q) , so by comparing degrees of both sides of $d'\eta_{r-1} = \xi' + d\eta'$, we get

$$d'\eta_{r-1} = d'\eta'_{r-1} + d''\eta'_r \text{ and } d''\eta'_{r-1} = 0.$$

If $d'\eta'_{r-1} \in \text{Im}d''$, then $d'\eta_{r-1} \in \text{Im}d''$ which contradicts to the fact that $\eta \in \mathcal{E}_r^{p-r, q+r-1}$. So $d'\eta'_{r-1} \notin \text{Im}d''$. Note that if η'_{r-1} is the leading term of a d -closed element τ , we may write $\tau = \eta'_{r-1} + \tau_r + \dots + \tau_k$ for some $k > r-1$ and each $\tau_i \in A^{p-r+i, q+r-1-i}$. Then comparing the degrees of $d'\tau = -d''\tau$, we get $d'\eta_{r-1} = -d''\tau_r$ which contradicts to the fact that $d'\eta_{r-1} \notin \text{Im}d''$.

From the above verification, we see that $\eta'_{r-1} \in \mathcal{E}_1^{p-1, q}$. Assume that $d\eta'_{r-1} \in Z_0^{p+1, q-1} + B_0^{p, q}$. Write $d\eta'_{r-1} = \gamma + d\sigma$ where $\gamma = \gamma_1 + \gamma_2 + \dots \in Z_0^{p+1, q-1}$, $\gamma_i \in A^{p+i, q-i}$, $\sigma = \sigma_0 + \sigma_1 + \dots \in B_0^{p, q}$ and $\sigma_i \in A^{p+i, q-1-i}$. Since the degree of $d\eta'_{r-1}$ is (p, q) , comparing the degrees of both sides of $d\eta'_{r-1} = \gamma + d\sigma$, we get $d\eta'_{r-1} = d''\sigma_0$ which contradicts to the fact that $\eta'_{r-1} \in \mathcal{E}_1^{p-1, q}$. Therefore $d\eta'_{r-1} \notin Z_0^{p+1, q-1} + B_0^{p, q}$ and hence $\beta_{p, q, 1}$ is not an isomorphism. \square

Theorem 2.11. Suppose that $(A = \oplus_{p, q \geq 0} A^{p, q}, d', d'')$ is a double complex and $r \geq 1$. The spectral sequence $\{E_r^{p, q}\}$ induced by A degenerates at E_r but not at E_{r-1} if and only if the following conditions hold:

- (1) $\mathcal{E}_k^{p, q} = \emptyset$ for all $p, q \in \mathbb{Z}, k \geq r$ and
- (2) $\mathcal{E}_{r-1}^{p, q} \neq \emptyset$ for some p, q .

Proof. Suppose that $\{E_r^{p, q}\}$ degenerates at E_r but not at E_{r-1} for some $r \geq 1$. By Proposition 2.7(2), $\alpha_{p, q, i}$ is an isomorphism for all $p, q \in \mathbb{Z}, i \geq r$. Then by Lemma 2.9, $\mathcal{E}_i^{p, q} = \emptyset$ for all $p, q \in \mathbb{Z}, i \geq r$. Since $d_{r-1} \neq 0$, by Proposition 2.7(1), there are some $p, q \in \mathbb{Z}$ such that $\beta_{p, q, r-1}$ is not an isomorphism. Then by Lemma 2.10, $\mathcal{E}_{r-1}^{p-r+1, q+r-2} \neq \emptyset$.

Conversely, suppose that (1) and (2) hold. By Lemma 2.10, $\beta_{p, q, k}$ is an isomorphism for all $p, q \in \mathbb{Z}, k \geq r$. Then by Proposition 2.7, $d_k = 0$ for $k \geq r$. For the case $r = 1$, by definition, $\mathcal{E}_0^{p, q} \neq \emptyset$ implies that $\beta_{p, q+1, 0}$ is not an isomorphism. And hence by Proposition 2.7, $d_0 \neq 0$. For the case $r \geq 2$, if $\beta_{p, q, r-1}$ is an isomorphism for all $p, q \in \mathbb{Z}$, by Proposition 2.7, $d_{r-1} = 0$. Then we have $d_k = 0$ for $k \geq r-1$. By the proof above, $\mathcal{E}_k^{p, q} = \emptyset$ for $k \geq r-1$. In particular, $\mathcal{E}_{r-1}^{p, q} = \emptyset$ for all $p, q \in \mathbb{Z}$ which contradicts to our assumption (2). Therefore there exist some p_0, q_0 such that $\beta_{p_0, q_0, r-1}$ is not an isomorphism. By Proposition 2.7, $d_{r-1} \neq 0$. \square

Definition 2.12. We say that a double complex (A, d', d'') satisfies the $d'd''$ -lemma at (p, q) if

$$\text{Im}d' \cap \ker d'' \cap A^{p, q} = \ker d' \cap \text{Im}d'' \cap A^{p, q} = \text{Im}d' d'' \cap A^{p, q}$$

and A satisfies the $d'd''$ -lemma if A satisfies the $d'd''$ -lemma at (p, q) for all (p, q) .

Now we can give a proof of the main result Theorem 1.1.

Proof. Note that by definition, $d'd''$ -lemma implies that $\text{Im}d' \cap \ker d'' \cap A^{p, q} = \text{Im}d' \cap \text{Im}d'' \cap A^{p, q}$ for all p, q . Since $\{E_r^{p, q}\}$ does not degenerate at E_0 , $\beta_{p, q, 0}$ is not an isomorphism for some p, q , hence by Lemma 2.10, $\mathcal{E}_0^{p, q-1} \neq \emptyset$. Assume that $\mathcal{E}_r^{p, q} \neq \emptyset$ for some $p, q \in \mathbb{Z}, r \geq 1$. Then there is $\alpha = \sum_{i=0}^{r-1} \alpha_i \in \mathcal{E}_r^{p, q}$ where $\alpha_i \in A^{p+i, q-i}$. From the condition $d\alpha = d'\alpha_{r-1}$, we have $d''\alpha_{r-1} = -d'\alpha_{r-2}$ and hence $d''d\alpha = -d'd''\alpha_{r-1} = 0$. So $d\alpha = d'\alpha_{r-1} \in (\text{Im}d' \cap \ker d'') \cap A^{p, q} = (\text{Im}d' \cap \text{Im}d'') \cap A^{p, q}$. But by the definition of $\mathcal{E}_r^{p, q}$, $d'\alpha_{r-1} \notin \text{Im}d''$ which leads to a contradiction. Therefore by Theorem 2.11, $\{E_r^{p, q}\}$ degenerates at E_1 . \square

In the following, we apply the main result to prove the E_1 -degeneration of a spectral sequence of bi-generalized Hermitian manifolds. We refer the reader to [G1, C] for generalized complex geometry, and to [CHT] for bi-generalized complex manifolds. We give a brief recall here. A bi-generalized complex structure on a smooth manifold M is a pair $(\mathcal{J}_1, \mathcal{J}_2)$ where $\mathcal{J}_1, \mathcal{J}_2$ are commuting generalized complex structures on M . A bi-generalized complex manifold is a smooth manifold M with a bi-generalized complex structure. A bi-generalized Hermitian manifold $(M, \mathcal{J}_1, \mathcal{J}_2, \mathbb{G})$ is an oriented bi-generalized complex manifold $(M, \mathcal{J}_1, \mathcal{J}_2)$ with a generalized metric \mathbb{G} which commutes with \mathcal{J}_1 and \mathcal{J}_2 . We define

$$U^{p,q} := U_1^p \cap U_2^q$$

where $U_1^p, U_2^q \subset \Gamma(\Lambda^* \mathbb{T}M \otimes \mathbb{C})$ are eigenspaces of $\mathcal{J}_1, \mathcal{J}_2$ associated to the eigenvalues ip and iq respectively and $\mathbb{T}M = TM \oplus T^*M$ is the generalized tangent space. It can be shown that the exterior derivative d is an operator from $U^{p,q}$ to $U^{p+1,q+1} \oplus U^{p+1,q-1} \oplus U^{p-1,q+1} \oplus U^{p-1,q-1}$ and we write

$$\delta_+ : U^{p,q} \rightarrow U^{p+1,q+1}, \delta_- : U^{p,q} \rightarrow U^{p+1,q-1}$$

for the projection of d into corresponding spaces.

Definition 2.13. *On a bi-generalized Hermitian manifold M , there is a double complex $\{(A, d', d'')\}$ given by*

$$A^{p,q} := U^{p+q,p-q}, d' = \delta_+, d'' = \delta_-$$

We call the spectral sequence $\{E_^{*,*}\}$ associated to this double complex the ∂_1 -Hodge-de Rham spectral sequence.*

By Theorem 1.1, we have the following result.

Theorem 2.14. *Suppose that M is a compact bi-generalized Hermitian manifold which satisfies the $\delta_+ \delta_-$ -lemma and has positive dimension. Then the ∂_1 -Hodge-de Rham spectral sequence degenerates at E_1 .*

Now we give a proof of the E_1 -degeneration of the ∂_1 -Hodge-de Rham spectral sequence.

Proof. Since $\bigoplus_{p,q} U^{p,q} = \Omega^*(M) \otimes \mathbb{C}$ (see [Ca07], pg 36) where $\Omega^*(M)$ is the collection of smooth forms on M , some $U^{p,q}$ is not empty. The space $U^{p,q}$ is a $C^\infty(M, \mathbb{C})$ -module where $C^\infty(M, \mathbb{C})$ is the ring of complex-valued smooth functions on M , and M has positive dimension, therefore $U^{p,q}$ is an infinite dimensional complex vector space. If δ_- is a zero map, we have $H_{\delta_-}^{p,q}(M) = U^{p,q}$ for all p, q . But M is compact, this contradicts to the fact that $H_{\delta_-}^{p,q}(M)$ is finite dimensional ([CHT, Theorem 2.14, Corollary 3.11]). Hence δ_- is not the zero map, and the spectral sequence does not degenerate at E_0 . Since we assume that M satisfies the $\delta_+ \delta_-$ -lemma, by Theorem 1.1, the spectral sequence degenerates at E_1 . \square

REFERENCES

- [C] G. Cavalcanti, New aspects of the dd^c -lemma, Oxford Univ. DPhil. thesis, arXiv:math/0501406v1[math.DG].
- [Ca07] G. Cavalcanti, Introduction to generalized complex geometry, impa, 26-Coloquio Brasileiro de Matematica, 2007.
- [CHT] T.W. Chen, C.I. Ho, and J.H. Teh, Aeppli and Bott-Chern cohomology for bigeneralized Hermitian manifolds and $d'd''$ -lemma, Journal of Geometry and Physics, 93(2015), 40-51.
- [DGMS] P. Deligne, P. Griffiths, J. Morgan, and D. Sullivan, *Real homotopy theory of Kähler manifolds*, Invent. Math., (3) 29 (1975), 245–274.
- [G1] M. Gualtieri, Generalized complex geometry, Ann. of Math., 174(2011), 75–123.

[M] J. McCleary, A user's Guide to spectral sequences, 2nd edition, Cambridge studies in advanced mathematics, Vol. 58, Cambridge University Press, 2001.

MATHEMATICS DEPARTMENT, NATIONAL TSING HUA UNIVERSITY, HSINCHU, TAIWAN
E-mail address: d937203@oz.nthu.edu.tw

MATHEMATICS DEPARTMENT, NATIONAL TSING HUA UNIVERSITY, NATIONAL CENTER OF THEORETICAL SCIENCES, MATHEMATICAL DIVISION, HSINCHU, TAIWAN
E-mail address: ciho@math.cts.nthu.edu.tw

MATHEMATICS DEPARTMENT, NATIONAL TSING HUA UNIVERSITY, HSINCHU, TAIWAN
E-mail address: jyhhaur@math.nthu.edu.tw